# datasheet NS Reagents



## **Anti-TREM2 Antibody**

Alternative Names: TREM-2, TREM2a, TREM2b, TREM2c, Trggering receptor expressed on myeloid cells 2,

Trggering receptor expressed on myeloid cells 2a, Triggering receptor expressed on

monocytes 2

Catalogue Number: AB21-10133-50ug

Size: 50 µg

## **Background Information**

Triggering receptor expressed on myeloid cells 2 (TREM2) is a cell surface transmembrane glycoprotein with V-immunoglobulin extracellular domains and cytoplasmic tails [1]. It is expressed in myeloid cells including dendritic cells, granulocytes, and tissue-specific macrophages such as osteoclasts. In the brain, TREM2 is only expressed by microglia and in the central nervous system expression of TREM2 varies with higher expression in the hippocampus, spinal cord and white matter.

TREM2 plays a complex role in neuroinflammation with expression of TREM2 being upregulated in pathological conditions such as Parkinson's disease, Amyotrophic lateral sclerosis (ALS), stroke, traumatic brain injury and Alzheimer's Disease [2]. TREM2 has also been shown to have a role in potential mechanisms linking urban air pollution to Alzheimer's Disease through its involvment in the regulation of neuroinflammation [3].

#### **Product Information**

Antibody Type: Polyclonal Host: Rabbit

Isotype: IgG Species Reactivity: Human, Mouse, Rat,

Dog, Pig, Cow, Horse

Immunogen: KLH conjugated synthetic peptide from human TREM2

Format: 50 μg in 50 μl PBS with 0.03% Proclin300, 50% glycerol, pH7.3.

Storage Conditions: 6 months: 4°C. Long-term storage: -20°C. Avoid multiple freeze and thaw cycles.

Applications: WB | ELISA | IHC-P | IHC-F | ICC | IF

WB 1:500-2000. ELISA 1:500-1000. IHC-P 1:100-500. IHC-F 1:100-500. ICC 1:100-500. IF 1:100-500.

#### **Additional Information**

Subcellular location: Cell membrane, Secreted MW: 25kDa (Intended as a general guide and does not

allow for all isoforms and species variations)

Gene ID 54209 Uniprot ID: Q9NZC2

tel: +44(0)1638 551500 | fax: +44(0)1638 590481 | tech@nktscientific.com | newmarketscientific.com

### **References**

[1] Bouchon A, Dietrich J, Colonna M. Cutting edge: inflammatory responses can be triggered by TREM-1, a novel receptor expressed on neutrophils and monocytes. J Immunol. 2000;164:4991–5.

[2] Gratuze, M., Leyns, C.E.G. & Holtzman, D.M. New insights into the role of TREM2 in Alzheimer's disease. Mol Neurodegeneration 13, 66 (2018).

[3] Hendrik J. Greve, Christen L. Mumaw, Evan J. Messenger, Prasada R. S. Kodavanti, Joyce L. Royland, Urmila P. Kodavanti and Michelle L. Block. Diesel exhaust impairs TREM2 to dysregulate neuroinflammation. J Neuroinflammation. 2020; 17: 351.

## **Images**

IHC of FFPE Rat brain TREM2: Cat No: AB21-10133 Antibody dilution: 1:200

